



Nonlinear Variational Inequalities and Implicit Variational Inequality of Ky Fan Type in H -Spaces

XIAN WU

Department of Mathematics, Yunnan Normal University
Kunming, Yunnan 650092, P.R. China

XIAN-ZHI YUAN

Department of Mathematics, The University of Queensland
Brisbane Qld 4072, Australia

(Received November 1998; revised and accepted May 1999)

Abstract—In this paper, the character of solution sets for nonlinear variational inequalities and the existence of a solution of implicit variational inequality of Ky Fan type in H -spaces are given.
© 1999 Elsevier Science Ltd. All rights reserved.

Keywords—L.c.-space, Locally convex H -space, Interval space, Variational inequality, Monotone function (mapping).

1. INTRODUCTION AND PRELIMINARIES

In 1977, Tarafdar [1] gave the following theorem.

THEOREM A. *Let E be a Hausdorff topological vector space, E^* the conjugate space of E and K a nonempty convex subset of E . Let $\langle \cdot, \cdot \rangle$ denote the pairing between E^* and E . If $T : K \rightarrow E^*$ is monotone and hemicontinuous, then a point $u_0 \in K$ is a solution of the variational inequality*

$$\langle T(u_0), v - u_0 \rangle \geq 0, \quad \forall v \in K$$

iff

$$\langle T(v), v - u_0 \rangle \geq 0, \quad \forall v \in K.$$

Using the theorem, Tarafdar [1] proved the existence of solutions of nonlinear variational inequality.

In 1991, Chang [2] generalized the above theorem as follows.

This work is supported by National Natural Science Foundation of China and Foundation of the Yunnan Sci. Tech. Commission and Foundation of Yunnan Educational Commission.

THEOREM B. Let X be a nonempty closed convex subset of a Hausdorff topological vector space E , $f : X \rightarrow (-\infty, +\infty]$ a lower semicontinuous function with $f \not\equiv +\infty$ and $\phi : X \times X \rightarrow R$ a monotone hemicontinuous (i.e., for each $y \in X$, $\phi(\cdot, y)$ is upper semicontinuous on each line-segment in X) function with $\phi(x, x) \geq 0$ for all $x \in X$. If the following conditions are fulfilled:

- (i) for each $x \in X$, $\phi(x, \cdot)$ is lower semicontinuous;
- (ii) for each $x \in X$, $f(y) + \phi(x, y)$ is a convex function in y ;

then there exists a point $\bar{x} \in X$ such that

$$f(y) + \phi(\bar{x}, y) \geq f(\bar{x}), \quad \forall y \in X,$$

iff

$$f(y) - \phi(y, \bar{x}) \geq f(\bar{x}), \quad \forall y \in X.$$

As an application of Theorem B, Chang [2] determined the character of solution sets for nonlinear variational inequalities in topological vector spaces.

In the present paper, our purpose is to establish a topological version of Theorems A and B and determine the character of solution sets for nonlinear variational inequalities and the existence of solution of implicit variational inequality of Ky Fan type in H -spaces.

We first give some definitions and notations.

Let X be a topological space and let $\mathcal{F}(X)$ be the family of all nonempty finite subsets of X . Let $\{\Gamma_A\}$ be a family of nonempty contractible subsets of X indexed by $A \in \mathcal{F}(X)$ such that $\Gamma_A \subset \Gamma_{A'}$ whenever $A \subset A'$. The pair $(X, \{\Gamma_A\})$ is called an H -space. Given an H -space $(X, \{\Gamma_A\})$, a nonempty subset D of X is called H -convex if $\Gamma_A \subset D$ for all $A \in \mathcal{F}(D)$; see [2–6].

An H -space $(X, \{\Gamma_A\})$ is called

- (1) a locally convex H -space if X is a uniform space and if there exists a base $\{V_i : i \in I\}$ for the uniform structure \mathcal{U} such that for each $i \in I$, $V_i(x) = \{y \in X : (y, x) \in V_i\}$ is H -convex for each $x \in X$ (see [7]);
- (2) an l.c.-space (see [4]) if X is a uniform space and if there exists a base $\{V_i : i \in I\}$ for the uniform structure such that for each $i \in I$, the set $\{x \in X : E \cap V_i[x] \neq \emptyset\}$ is H -convex whenever E is H -convex, where $V_i[x] = \{y \in X : (x, y) \in V_i\}$.

REMARK. The concept of an l.c.-space is different from a locally convex H -space. But an l.c.-space $(X, \{\Gamma_A\})$ with $\Gamma_{\{x\}} = \{x\}$ for all $x \in X$ must be a locally convex H -space. A nonempty convex subset X of a locally convex topological vector space is an l.c.-space with $\Gamma_A = \text{co } A$ for all $A \in \mathcal{F}(X)$, and hence, $(X, \{\text{co } A\})$ must be a locally convex H -space.

Let X be an H -space and $f : X \rightarrow R \cup \{+\infty\}$ a function. f is called H -quasiconvex if for each $r \in R$, the set $\{x \in X : f(x) < r\}$ is H -convex; f is called strongly H -quasiconvex if f is H -quasiconvex and for each $r \in R$, if $f(y_1) < r$ and $f(y_2) \leq r$ implies that $f(y) < r$ for all $y \in \Gamma_{\{y_1, y_2\}} \setminus \{y_1, y_2\}$; f is called strictly H -quasiconvex if f is H -quasiconvex and for any $y_1, y_2 \in X$ with $y_1 \neq y_2$, $f(y) < \max\{f(y_1), f(y_2)\}$ for all $y \in \Gamma_{\{y_1, y_2\}} \setminus \{y_1, y_2\}$.

If a topological space X is endowed with a mapping

$$[\cdot, \cdot] : X \times X \rightarrow \{\text{connected subsets of } X\},$$

such that $x_1, x_2 \in [x_1, x_2] = [x_2, x_1]$ for all $x_1, x_2 \in X$, then $(X, [\cdot, \cdot])$ is called an interval space, $[x_1, x_2]$ is called an interval. Obviously, an H -space $(X, \{\Gamma_A\})$ with $\Gamma_{\{x\}} = \{x\}$ ($\forall x \in X$) is an interval space and $[x_1, x_2] = \Gamma_{\{x_1, x_2\}}$ ($x_1, x_2 \in X$).

Given an interval space $(X, [\cdot, \cdot])$, use w -convex consistently in the whole paper. $f : X \rightarrow R \cup \{+\infty\}$ is called w -convex iff for each $r \in R$, if $f(y_1) < r$ and $f(y_2) \leq r$ implies that $f(y) < r$ for all $y \in [y_1, y_2] \setminus \{y_1, y_2\}$.

Let X be a topological space. We denote by 2^X the family of all subsets of X . If $A \subset X$ we shall denote by $\text{cl}(A)$ the closure of A . If X is a topological vector space, we shall denote the convex hull of A by $\text{co } A$.

A topological space is called acyclic if all of its reduced Čech homology groups over rationals vanish. In particular, any contractible space is acyclic, and thus, any convex or star-shaped set is acyclic.

Let X, Y be two topological spaces, $T : X \longrightarrow 2^Y$ a multivalued mapping and $\phi : X \times X \longrightarrow R$ a function.

- (1) T is called upper semicontinuous if for each $x \in X$ and each open set $V \subset Y$ with $T(x) \subset V$, there exists an open neighborhood U of x such that $T(z) \subset V$ for each $z \in U$.
- (2) ϕ is called monotone iff

$$\phi(x, y) + \phi(y, x) \leq 0,$$

for all $(x, y) \in X \times X$.

- (3) The multivalued mapping $\text{cl}T : X \longrightarrow 2^Y$ is defined by

$$\text{cl}T(x) = \text{cl}[T(x)], \quad \forall x \in X.$$

Let X be a nonempty convex subset of a Hausdorff topological vector space E , E^* the conjugate space of E ($\langle \cdot, \cdot \rangle$ denotes the pair in duality), and $A : X \longrightarrow E^*$ a mapping.

- (1) A is called monotone if $\text{Re}\langle A(x_1) - A(x_2), x_1 - x_2 \rangle \geq 0$ for all $x_1, x_2 \in X$.
- (2) A is called hemicontinuous if A is continuous from the line-segments in X to the weak topology of E^* .

REMARK. Let $\phi(x, y) = \langle A(x), y - x \rangle$, $x, y \in X$. If A is monotone and hemicontinuous, then ϕ is also monotone and hemicontinuous.

2. A TOPOLOGICAL TOOL

THEOREM 1. Let $(X, [\cdot, \cdot])$ be a Hausdorff interval space, $f : X \rightarrow (-\infty, +\infty]$ a lower semicontinuous function with $f \not\equiv +\infty$, and $\phi : X \times X \longrightarrow R$ a monotone function with $\phi(x, x) \geq 0$ for all $x \in X$. If the following conditions are fulfilled:

- (i) for each $y \in X$, $\phi(\cdot, y)$ is upper semicontinuous on each interval;
- (ii) for each $x \in X$, $f(y) + \phi(x, y)$ is w -convex in y ;

then there exists a point $\bar{x} \in X$ such that

$$f(y) + \phi(\bar{x}, y) \geq f(\bar{x}), \quad \forall y \in X,$$

iff

$$f(y) - \phi(y, \bar{x}) \geq f(\bar{x}), \quad \forall y \in X.$$

PROOF. If there exists a point $\bar{x} \in X$ such that

$$f(y) + \phi(\bar{x}, y) \geq f(\bar{x}), \quad \forall y \in X,$$

then

$$f(y) - \phi(y, \bar{x}) \geq f(y) + \phi(\bar{x}, y) \geq f(\bar{x}),$$

for all $y \in X$ since ϕ is monotone. Contrarily, if there exists a point $\bar{x} \in X$ such that

$$f(y) - \phi(y, \bar{x}) \geq f(\bar{x}), \quad \forall y \in X, \tag{1}$$

but, there is a point $\bar{y} \in X$ such that

$$f(\bar{y}) + \phi(\bar{x}, \bar{y}) < f(\bar{x}),$$

then

$$\bar{x} \in U := \{z \in [\bar{x}, \bar{y}] : f(z) - \phi(z, \bar{y}) > f(\bar{y})\}.$$

Obviously, $\bar{y} \notin U$ and U is open in $[\bar{x}, \bar{y}]$ since both f and $-\phi(\cdot, \bar{y})$ are lower semicontinuous on $[\bar{x}, \bar{y}]$.

If $([\bar{x}, \bar{y}] \setminus \{\bar{x}, \bar{y}\}) \cap U = \emptyset$, then $U = \{\bar{x}\}$ is closed in $[\bar{x}, \bar{y}]$. It contradicts the connectivity of $[\bar{x}, \bar{y}]$. Hence, there exists a point $\bar{z} \in ([\bar{x}, \bar{y}] \setminus \{\bar{x}, \bar{y}\}) \cap U$, i.e., $\bar{z} \in ([\bar{x}, \bar{y}] \setminus \{\bar{x}, \bar{y}\})$ and

$$f(\bar{y}) + \phi(\bar{z}, \bar{y}) < f(\bar{z}). \quad (2)$$

Taking $y = \bar{z}$ in (1) we have

$$f(\bar{x}) + \phi(\bar{z}, \bar{x}) \leq f(\bar{z}). \quad (3)$$

By (ii) together with (2) and (3), we have

$$f(x) + \phi(\bar{z}, x) < f(\bar{z}),$$

for all $x \in [\bar{x}, \bar{y}] \setminus \{\bar{x}, \bar{y}\}$, and hence, $\phi(\bar{z}, \bar{z}) < 0$. It contradicts that $\phi(x, x) \geq 0$ for all $x \in X$. Hence,

$$f(y) + \phi(\bar{x}, y) \geq f(\bar{x}), \quad \forall y \in X.$$

This completes the proof.

COROLLARY 2. *Let X be a nonempty convex subset of a Hausdorff topological vector space E , $f : X \rightarrow (-\infty, +\infty]$ a lower semicontinuous function with $f \not\equiv +\infty$ and $\phi : X \times X \rightarrow R$ a monotone function with $\phi(x, x) \geq 0$ for all $x \in X$. If the following conditions are fulfilled:*

- (i) *for each $y \in X$, $\phi(\cdot, y)$ is upper semicontinuous on each line-segment;*
- (ii) *for each $x \in X$, $f(y) + \phi(x, y)$ is a convex function in y ;*

then there exists a point $\bar{x} \in X$ such that

$$f(y) + \phi(\bar{x}, y) \geq f(\bar{x}), \quad \forall y \in X,$$

iff

$$f(y) - \phi(y, \bar{x}) \geq f(\bar{x}), \quad \forall y \in X.$$

PROOF. For any $x_1, x_2 \in X$, let $[x_1, x_2] = \text{co}\{x_1, x_2\}$. Then X is an interval space. Moreover, by (ii) we can prove that $f(y) + \phi(x, y)$ is w -convex in y for each $x \in X$. Consequently, the conclusion follows from Theorem 1.

REMARK. Corollary 2 shows that X need not be closed and $\phi(x, \cdot)$ need not be lower semicontinuous in [2, Theorem 3.9.2] (i.e., Theorem B).

3. SOLUTION SET FOR NONLINEAR VARIATIONAL INEQUALITIES

THEOREM 3. *Let $(X, \{\Gamma_A\})$ be a Hausdorff H -space with $\Gamma_{\{x\}} = \{x\}$ for all $x \in X$, $f : X \rightarrow R \cup \{+\infty\}$ a lower semicontinuous function with $f \not\equiv +\infty$ and $\phi : X \times X \rightarrow R$ a monotone function with $\phi(x, x) \geq 0$ for all $x \in X$. If the following conditions are fulfilled:*

- (i) *there exist a point $x_0 \in X$ and a compact subset K of X such that*

$$f(x) > f(x_0) + \phi(x, x_0), \quad \forall x \in X \setminus K;$$

- (ii) *for each $x \in X$, $f(y) + \phi(x, y)$ is strongly H -quasiconvex in y ;*
- (iii) *for each $y \in X$, $\phi(\cdot, y)$ is upper semicontinuous on $\Gamma_{\{x_1, x_2\}}$ for each $(x_1, x_2) \in X \times X$;*
- (iv) *for each $x \in X$, $\phi(x, \cdot)$ is lower semicontinuous;*

then the solution set of the variational inequality

$$\phi(\bar{x}, y) \geq f(\bar{x}) - f(y), \quad \forall y \in X, \quad (*)$$

is a nonempty compact H -convex subset of K .

If (ii) is replaced by the following:

(ii)' for each $x \in X$, $f(y) + \phi(x, y)$ is strictly H -quasiconvex in y ,

then the variational inequality $(*)$ has a unique solution \bar{x} and $\bar{x} \in K$.

PROOF. For any $x_1, x_2 \in X$, let $[x_1, x_2] = \Gamma_{\{x_1, x_2\}}$. Then X is an interval space. For each $y \in X$, let

$$M(y) = \{x \in X : f(y) + \phi(x, y) \geq f(x)\}, \quad H(y) = \{x \in X : f(y) - \phi(y, x) \geq f(x)\}.$$

Then $H(y)$ is closed and H -convex by (ii), (iv), and the lower semicontinuity of f . Let

$$S = \bigcap_{y \in X} M(y).$$

Then S is the set of all solutions of the variational inequality $(*)$. By Theorem 1 we know that $S = \bigcap_{y \in X} H(y)$.

For each finite subset $\{y_1, y_2, \dots, y_n\} \subset X$, if $x \notin \bigcup_{i=1}^n M(y_i)$, then $f(y_i) + \phi(x, y_i) < f(x)$ for all $i \in \{1, 2, \dots, n\}$. Since $f(y) + \phi(x, y)$ is H -quasiconvex in y ,

$$\Gamma_{\{y_1, y_2, \dots, y_n\}} \subset \{y \in X : f(y) + \phi(x, y) < f(x)\}$$

so that $x \notin \Gamma_{\{y_1, y_2, \dots, y_n\}}$ because $\phi(x, x) \geq 0$. This shows that $M : X \rightarrow 2^X$ is an H -KKM mapping so that $\text{cl } M : X \rightarrow 2^X$ is also an H -KKM mapping. Moreover, by (i) we know that $M(x_0) \subset K$, and hence, $\text{cl } [M(x_0)] \subset K$ is compact. By virtue of the H -KKM theorem (see also [2-4]), we know that

$$\bigcap_{y \in X} \text{cl } [M(y)] \neq \emptyset.$$

Noting that $\text{cl } [M(y)] \subset H(y)$ for all $y \in X$, we know that

$$S = \bigcap_{y \in X} H(y) \neq \emptyset.$$

Consequently, S is a nonempty compact H -convex subset of K since

$$\bigcap_{y \in X} H(y) = S \subset \text{cl } S \subset \text{cl } [M(x_0)] \subset K$$

and $H(y)$ is closed and H -convex for all $y \in X$.

If (ii) is replaced by (ii)', then for any $x_1, x_2 \in S$, we have

$$f(y) + \phi(x_i, y) \geq f(x_i), \quad \forall y \in X, \quad (4)$$

$i = 1, 2$. Since ϕ is monotone and $\phi(x, x) \geq 0$, $\phi(x, x) = 0$ for all $x \in X$. If $x_1 \neq x_2$, taking $y \in \Gamma_{\{x_1, x_2\}} \setminus \{x_1, x_2\}$ in (4) we have

$$f(x_1) \leq f(y) + \phi(x_1, y) < f(x_2) + \phi(x_1, x_2)$$

and

$$f(x_2) \leq f(y) + \phi(x_2, y) < f(x_1) + \phi(x_2, x_1)$$

by (ii)'.

Consequently,

$$\phi(x_1, x_2) > f(x_1) - f(x_2)$$

and

$$\phi(x_2, x_1) > f(x_2) - f(x_1),$$

and hence,

$$\phi(x_1, x_2) + \phi(x_2, x_1) > 0.$$

It contradicts the monotoneity of ϕ . Therefore, S contains a unique element; i.e., the variational inequality $(*)$ has a unique solution \bar{x} and $\bar{x} \in K$.

COROLLARY 4. *Let X be a nonempty closed convex subset of a Hausdorff topological vector space E , $f : X \rightarrow R$ a lower semicontinuous function and $\phi : X \times X \rightarrow R$ a monotone function with $\phi(x, x) \geq 0$ for all $x \in X$. If the following conditions are fulfilled:*

- (i) *there exist a compact subset K of E and a point $x_0 \in X$ such that*

$$f(x) > \phi(x, x_0) + f(x_0), \quad \forall x \in X \setminus K;$$

- (ii) *for each $x \in X$, $f(y) + \phi(x, y)$ is a convex function in y ;*
 (iii) *for each $y \in X$, $\phi(\cdot, y)$ is upper semicontinuous on each line-segment in X ;*
 (iv) *for each $x \in X$, $\phi(x, \cdot)$ is lower semicontinuous;*

then the solution set of the variational inequality

$$\phi(x, y) \geq f(x) - f(y), \quad \forall y \in X, \quad (*)$$

is a nonempty compact convex subset of $X \cap K$.

If (ii) is replaced by the following:

- (ii)' *for each $x \in X$, $f(y) + \phi(x, y)$ is strictly convex in y ,*

then the variational inequality () has a unique solution \bar{x} and $\bar{x} \in X \cap K$.*

PROOF. For each finite subset A of X , let $\Gamma_A = \text{co } A$. Then $(X, \{\text{co } A\})$ is a Hausdorff H -space. Since $\phi(x, x) \geq 0$ for all $x \in X$, $x_0 \in X \cap K$ by (i). Since again X is closed in E , $X \cap K$ is a nonempty compact subset of X . Note that $X \setminus K = X \setminus (X \cap K)$. By (i) we know that

$$f(x) > \phi(x, x_0) + f(x_0), \quad \forall x \in X \setminus (X \cap K).$$

Consequently, the corollary follows from Theorem 3.

REMARK. Corollary 4 shows that $\phi(\cdot, y)$ need not be upper semicontinuous in [2, Theorem 3.9.3].

COROLLARY 5. *Let X be a nonempty compact convex subset of a Hausdorff topological vector space E and E^* the conjugate space of E ($\langle \cdot, \cdot \rangle$ denotes the pair in duality). Let $A : X \rightarrow E^*$ be monotone and hemicontinuous. Then the solution set of the variational inequality*

$$\text{Re}\langle Ax, y - x \rangle \geq 0, \quad \forall y \in X, \quad (*)$$

is a nonempty compact convex subset of X .

PROOF. For each $(x, y) \in X \times X$, let $\phi(x, y) = \text{Re}\langle Ax, y - x \rangle$. Then $\phi : X \times X \rightarrow R$ satisfies all conditions of Corollary 4. Consequently, the corollary follows from Corollary 4.

REMARK. Corollary 5 is finer than the corollary in [1].

4. KY FAN TYPE IMPLICIT VARIATIONAL INEQUALITY IN H -SPACES

First, we give a lemma, which appeared in [7].

LEMMA 6. *Let $(X, \{\Gamma_A\})$ be a compact Hausdorff locally convex H -space and $T : X \rightarrow 2^X$ an upper semicontinuous multivalued mapping with closed acyclic values. Then there exists a point $x_0 \in X$ such that $x_0 \in T(x_0)$.*

THEOREM 7. *Let $(X, \{\Gamma_A\})$ be a compact Hausdorff l.c.-space with $\Gamma_{\{x\}} = \{x\}$ for all $x \in X$, $g : X \times X \rightarrow R \cup \{+\infty\}$ a lower semicontinuous function and $\psi : X \times X \times X \rightarrow R$ a function with $\psi(z, x, x) \geq 0$ for all $(z, x) \in X \times X$. If the following conditions are fulfilled:*

- (i) *for each $z \in X$, there exists a point $x \in X$ such that $g(z, x) < +\infty$;*
 (ii) *for each $z \in X$, $\phi(x, y) = \psi(z, x, y)$ is monotone;*
 (iii) *for each $x \in X$, $g(\cdot, x)$ is upper semicontinuous;*
 (iv) *for each $(z, x) \in X \times X$, $g(z, y) + \psi(z, x, y)$ is strictly H -quasiconvex in y ;*
 (v) *$\psi(z, \cdot, y)$ is upper semicontinuous on each $\Gamma_{\{x_1, x_1\}}(x_1, x_1 \in X)$ and $\psi(\cdot, x, \cdot)$ is lower semicontinuous;*

then there exists a point $\bar{x} \in X$ such that

$$g(\bar{x}, y) + \psi(\bar{x}, \bar{x}, y) \geq g(\bar{x}, \bar{x}), \quad \forall y \in X.$$

PROOF. For any $x_1, x_2 \in X$, let $[x_1, x_2] = \Gamma_{\{x_1, x_2\}}$. Then $(X, [\cdot, \cdot])$ is a compact Hausdorff interval space. For each fixed $z \in X$, let $f(x) = g(z, x)$ for all $x \in X$. Then f, ϕ satisfies all conditions of Theorem 1. By Theorem 1, $\bar{x} \in X$ is a solution of the variational inequality

$$f(y) + \phi(x, y) \geq f(x), \quad \forall y \in X, \quad (5)$$

iff $\bar{x} \in X$ is a solution of the variational inequality

$$f(y) - \phi(y, x) \geq f(x), \quad \forall y \in X. \quad (6)$$

Note that f, ϕ satisfy all conditions of the second part of Theorem 3. By virtue of Theorem 3 we know that the variational inequality (5) has unique solution $s(z)$, and hence, the variational inequality (6) has unique solution $s(z)$.

Let $\{(z_\alpha, s(z_\alpha)) : \alpha \in I\}$ be a net converging to (z, x) . Then for each $y \in X$, we have

$$g(z_\alpha, y) - \psi(z_\alpha, y, s(z_\alpha)) \geq g(z_\alpha, s(z_\alpha))$$

for all $\alpha \in I$. By (iii), (v), and the lower semicontinuity of g , we know that

$$\begin{aligned} g(z, x) &\leq \underline{\lim}_\alpha g(z_\alpha, s(z_\alpha)) \\ &\leq \underline{\lim}_\alpha [g(z_\alpha, y) - \psi(z_\alpha, y, s(z_\alpha))] \\ &\leq \overline{\lim}_\alpha [g(z_\alpha, y) - \psi(z_\alpha, y, s(z_\alpha))] \\ &\leq g(z, y) - \psi(z, y, x), \end{aligned}$$

for all $y \in X$. This shows that $x = s(z)$, and hence, the graph of s is closed and $s : X \rightarrow X$ is a single-valued upper semicontinuous mapping.

Note that $(X, \{\Gamma_A\})$ is also a compact Hausdorff locally convex H -space. By virtue of [7, Theorem 1], i.e., Lemma 6, there exists a point $\bar{x} \in X$ such that $\bar{x} = s(\bar{x})$, i.e.,

$$g(\bar{x}, y) + \psi(\bar{x}, \bar{x}, y) \geq g(\bar{x}, \bar{x}), \quad \forall y \in X.$$

This completes the proof.

COROLLARY 8. Let $(X, \{\Gamma_A\})$ be a compact Hausdorff l.c.-space with $\Gamma_{\{x\}} = \{x\}$ for all $x \in X$ and $\psi : X \times X \times X \rightarrow R$ a function with $\psi(z, x, x) \geq 0$ for all $(z, x) \in X \times X$. If the following conditions are fulfilled:

- (i) for each $z \in X$, $\phi(x, y) = \psi(z, x, y)$ is monotone;
- (ii) for each $(z, x) \in X \times X$, $\psi(z, x, y)$ is strictly H -quasiconvex in y ;
- (iii) $\psi(z, \cdot, y)$ is upper semicontinuous on each $\Gamma_{\{x_1, x_1\}}(x_1, x_1 \in X)$ and $\psi(\cdot, x, \cdot)$ is lower semicontinuous;

then there exists a point $\bar{x} \in X$ such that

$$\psi(\bar{x}, \bar{x}, y) \geq 0, \quad \forall y \in X.$$

PROOF. Take $g \equiv 0$ in Theorem 7. Then the conclusion follows from Theorem 7.

REMARK. Corollary 8 extends Ky Fan's theorem for implicit variational inequalities (see also [2, Theorem 6.7.1]) to H -spaces. Theorem 7 improves [2, Theorem 6.6.1].

REFERENCES

1. E. Tarafdar, On nonlinear variational inequalities, *Proc. Amer. Math. Soc.* **67** (1), 95–98, (1977).
2. S.S. Chang, *Variational Inequality and Complementarity Problem Theory and Applications*, Shanghai Scientific and Technological Literature Publishing, (1991).
3. C. Bardaro and R. Ceppitelli, Some further generalizations of Knaster-Kuratowski-Mazurkiewicz theorem and minimax inequalities, *J. Math. Anal. Appl.* **132**, 484–490, (1988).
4. S.S. Chang and Y.H. Ma, Generalized KKM theorem on H -spaces with applications, *J. Math. Anal. Appl.* **163**, 406–421, (1992).
5. C.D. Horvath, Contractibility and generalized convexity, *J. Math. Anal. Appl.* **156**, 341–357, (1991).
6. C.D. Horvath, *Some Results on Multivalued Mappings and Inequalities Without Convexity, Nonlinear and Convex Analysis*, (Edited by B.L. Lin and Simons), pp. 99–106, Marcel Dekker, New York, (1987).
7. X. Wu, Kakutani-Fan-Glicksberg fixed point theory with applications in H -spaces, *J. Math. Anal. Appl.* (to appear).
8. S. Kum, A generalization of quasi-variational inequalities, *J. Math. Anal. Appl.* **182**, 158–164, (1994).